

New Developments on Turán's Extremal Problems for Polynomials

Gradimir V. Milovanović

Department of Mathematics, University of Niš
Faculty of Electronic Engineering, P.O. Box 73, 18000 Niš, Yugoslavia

Themistocles M. Rassias

National Technical University of Athens
Department of Mathematics, Zografou Campus, 15780 Athens, Greece

Abstract

In this paper we give an account of L^r inequalities of Turán type for algebraic polynomials, mainly initiated and studied by the late Professor Arun K. Varma. This paper could be comprehended as a continuation of our previous survey paper [8].

1 Introduction

Let $\mathcal{P}_n$ be the set of all algebraic polynomials of degree at most n and let W_n be some of its subsets. For a given norm $\|\cdot\|$ we consider extremal problems

$$B_{n,m} = \inf_{P \in W_n} \frac{\|P^{(m)}\|}{\|P\|} \quad (1 \leq m \leq n).$$

In comparing with inequalities of Markov's type (cf. Milovanović, Mitrinović, Rassias [9, Chap. 6]), here we have opposite inequalities which are known as *inequalities of Turán type*.

Turán [11] proved the following inequality for polynomials $P \in \mathcal{P}_n$ having all their zeros in $[-1, 1]$,

$$\|P'\|_\infty > \frac{\sqrt{n}}{6} \|P\|_\infty, \quad (1)$$

taking the uniform norm $\|f\|_\infty = \max_{-1 \leq t \leq 1} |f(t)|$. The constant $\sqrt{n}/6$ is not the best possible.

Turán's inequality (1) has been generalized and extended in several different ways.

Firstly, inequality (1) was sharpened by Erőd [6], who obtained

$$\|P'\|_\infty \geq B_n \|P\|_\infty, \quad (2)$$

where $B_2 = 1$, $B_3 = 3/2$, and

$$B_{2k} = \frac{2k}{\sqrt{2k-1}} \left(1 - \frac{1}{2k-1}\right)^{k-1},$$

$$B_{2k+1} = \frac{(2k+1)^2}{2k\sqrt{2k+2}} \left(1 - \frac{\sqrt{2k+2}}{2k}\right)^{k-1} \left(1 + \frac{1}{\sqrt{2k+2}}\right)^k,$$

for $k = 2, 3, \dots$.

Exactly, equality in (2) is attained for $P(t) = (1-t)^n$, if $n = 1, 2, 3$, and for $P(t) = (1-t)^{n-[n/2]}(1+t)^{[n/2]}$, if $n \geq 4$.

Let W_n be the set of all algebraic polynomials of degree n whose zeros are all real and lie inside $[-1, 1]$. The corresponding inequality for the second derivative of such polynomials was investigated by Babenko and Pichugov [2].

If $P \in W_n$, $n \geq 2$, they proved that

$$\|P''\|_\infty \geq B_{n,2} \|P\|_\infty, \quad (3)$$

where $B_{n,2} = \min\{n, (n-1)n/4\}$.

If $n = 2, 3, 4, 5$, then $B_{n,2} = (n-1)n/4$, and equality in (3) is attained only for polynomials of the form $P(t) = C(1 \pm t)^n$, where C is an arbitrary real constant different from zero.

In the case $n \geq 6$, they found that $B_{n,2} = n$, and for $n = 2m$ equality in (3) holds only for polynomials of the form $P(t) = C(1-t^2)^m$, where C is an arbitrary real constant different from zero.

An analogue in L^2 norm for algebraic polynomials was considered firstly by Professor A. K. Varma [15]. Taking $\|f\|_2^2 = \int_{-1}^1 f(t)^2 dt$ he proved:

Theorem 1 *If $P \in W_n$ we have*

$$\|P'\|_2^2 \geq \frac{n}{2} \|P\|_2^2. \quad (4)$$

This result is best possible in the sense that there exists a polynomial P_0 of degree n having all zeros inside $[-1, 1]$ and for which

$$\|P'_0\|_2^2 = \left(\frac{n}{2} + \frac{3}{4} + \frac{3}{4(n-1)}\right) \|P_0\|_2^2, \quad n > 1.$$

The proof of this theorem was based on the following inequality

$$\| \sqrt{1-t^2} P' \|_2^2 \geq \frac{n}{2} \|P\|_2^2 \quad (P \in W_n),$$

which becomes an equality only for $P(t) = C(1+t)^p(1-t)^q$, $p+q=n$, where C is an arbitrary non-zero constant.

In this survey we give an account of L^r ($r \geq 1$) inequalities of Turán type.

2 Turán Type Inequalities in L^2 Norm

In [16] Professor Varma gave a more precise form of (4).

Theorem 2 Let $\|f\|_2^2 = \int_{-1}^1 f(t)^2 dt$, $P \in W_n$ and $P(1) = P(-1) = 0$. Then we have

$$\|P'\|_2^2 \geq \left(\frac{n}{2} + \frac{3}{4} + \frac{3}{4(n-1)} \right) \|P\|_2^2, \quad (5)$$

with equality for $P(t) = (1-t^2)^m$, $n = 2m$.

Taking the norm $\|f\|_2^2 = \int_{-1}^1 (1-t^2)f(t)^2 dt$, in 1979 Varma [17] proved the following result:

Theorem 3 For $P \in W_n$ and $n \geq 2$ we have

$$\|P'\|_2^2 \geq \left(\frac{n}{2} + \frac{1}{4} - \frac{1}{4(n+1)} \right) \|P\|_2^2,$$

with equality for $P(t) = (1-t^2)^m$, $n = 2m$.

Later Varma [18] proved an improvement of one of his earlier results.

Theorem 4 Let $\|f\|_2^2 = \int_{-1}^1 f(t)^2 dt$. If $P \in W_n$ and $n = 2m$; then

$$\|P'\|_2^2 \geq \left(\frac{n}{2} + \frac{3}{4} + \frac{3}{4(n-1)} \right) \|P\|_2^2, \quad (6)$$

where equality holds if and only if $P(t) = (1-t^2)^m$. Moreover, if $n = 2m-1$, then

$$\|P'\|_2^2 \geq \left(\frac{n}{2} + \frac{3}{4} + \frac{5}{4(n-2)} \right) \|P\|_2^2, \quad n \geq 3, \quad (7)$$

where equality holds if and only if $P(t) = (1-t)^{m-1}(1+t)^m$ or $P(t) = (1-t)^m(1+t)^{m-1}$.

This result is an improvement of Theorem 2 in two respects. First, the condition $P(1) = P(-1) = 0$ is not necessary for (5) to hold. Secondly, here there exist precise bounds for n even and also for n odd as mentioned in (6) and (7).

In the same norm, Varma [18] also proved:

Theorem 5 *Let $P \in W_n$, subject to the condition $P(1) = 1$; then*

$$\|P'\|_2^2 \geq \frac{n}{4} + \frac{1}{8} + \frac{1}{8(2n-1)}, \quad n \geq 1,$$

where equality holds for $P(t) = ((1+t)/2)^n$.

This inequality is an improvement over $\|P'\|_2^2 > n/4$, given by Szabados and Varma [10].

The corresponding inequality for polynomials $P \in W_n$ in L^r norm, defined on $(-1, 1)$ by $\|f\|_r = \left(\int_{-1}^1 |f(t)|^r dt \right)^{1/r}$, was considered by Zhou [20].

Theorem 6 *If $P \in W_n$, then for $1 \leq r \leq +\infty$,*

$$\|P'\|_r \geq C\sqrt{n} \|P\|_r,$$

where C is a positive absolute constant.

A similar result for $0 < r < 1$ was obtained also by Zhou [21]. Recently, Zhou [22] proved the following results:

Theorem 7 *If $P \in W_n$, then for $1 \leq r \leq s \leq +\infty$,*

$$\|P'\|_r \geq Cn^{1/2-1/(2r)+1/(2s)} \|P\|_s,$$

where C is a positive absolute constant.

The example $P(t) = (1 - t^2)^{[n/2]}$ in the previous theorem shows that the order $n^{1/2-1/(2r)+1/(2s)}$ cannot be improved.

Theorem 8 *Let $1 \leq r \leq s \leq +\infty$ and P be a polynomial of degree n with only real zeros. If at most k zeros of P lie outside the interval $[-1, 1]$, then*

$$\|P'\|_r \geq C_k n^{1/2-1/(2r)+1/(2s)} \|P\|_s,$$

where C_k is a positive constant depending only upon k .

After Professor Varma's death, the following result [14] has appeared:

Theorem 9 Let $\|f\|^2 = \int_{-1}^1 (1-t^2)^\alpha f(t)^2 dt$, $P \in W_n$ ($n \geq 2$) and $\alpha > 1$ real. Then we have ($n = 2m$)

$$\|P'\|^2 \geq \frac{n^2(2n+2\alpha+1)}{4(n+\alpha-1)(n+\alpha)} \|P\|^2,$$

with equality if and only if $P(t) = c(1-t^2)^m$. If $P(\pm 1) = 0$, then the previous inequality remains valid for $\alpha > -1$.

This result was proved earlier by Varma for the cases $\alpha = 0$ and $\alpha = 1$. In the same paper [14], Underhill and Varma investigated the corresponding inequality in L^4 norm for $\alpha = 3$:

Theorem 10 Let $\|f\|_4^4 = \int_{-1}^1 (1-t^2)^3 f(t)^4 dt$ and $P \in W_n$. Then we have ($n = 2m$)

$$\|P'\|_4^4 \geq \frac{3n^3(4n+7)(4n+5)}{4(4n+6)(4n+4)(4n+2)} \|P\|_4^4,$$

with equality if and only if $P(t) = c(1-t^2)^m$.

Also, they considered the cases when $\alpha = 1$ and $\alpha = 2$, as well as an inequality in L^r norm, when $r \geq 2$ is even. In [19] Varma proved:

Theorem 11 Let $P \in W_n$, subject to the condition $P(1) = 0$. Then, for $r \geq 1$, we have

$$\int_{-1}^1 |P'(t)|^r dt \geq \frac{n^r}{2^{r-1}((n-1)r+1)},$$

with equality if and only if $P(t) = ((1+t)/2)^n$.

3 Bojanov's Solution

More general results on Turán type inequalities were obtained by Bojanov [3]. Introducing the notations

$$p_{n,k}(t) = (-1)^{n-k} \frac{n^n}{2^n k^k (n-k)^{n-k}} (t+1)^k (t-1)^{n-k},$$

for $k = 0, 1, \dots, n$ ($n \in \mathbb{N}$), Bojanov [3] proved the following results:

Theorem 12 Let $x \mapsto \varphi(x)$ be any continuously differentiable, strictly increasing convex function in $[0, +\infty)$. Then for every $n \in \mathbb{N}$ and $m \in \{1, \dots, n\}$, there exists a constant $A_{n,m} > 0$ such that

$$\int_{-1}^1 \varphi(|P^{(m)}(t)|) dt \geq A_{n,m} \|P\|_\infty \quad (P \in W_n).$$

Moreover,

$$A_{n,m} = \min_{0 \leq k \leq n} \left\{ \int_{-1}^1 \varphi(|p_{n,k}^{(m)}(t)|) dt \right\}$$

and this is the exact constant.

Theorem 13 For any given n and m , there exists a constant $B_{n,m}$ such that

$$\|P^{(m)}\|_{\infty} \geq B_{n,m} \|P\|_{\infty} \quad (P \in W_n).$$

Moreover,

$$B_{n,m} = \min_{0 \leq k \leq n} \{\|p_{n,k}^{(m)}\|_{\infty}\}.$$

Using this theorem one could get the exact previous result of Erőd [6] and Babenko and Pichugov [2], treating the case $m = 1$ and $m = 2$, respectively. In the first case we have that

$$B_{n,1} = \|p'_{n,k}\|_{\infty} \quad \text{for } k = \left\lfloor \frac{n}{2} \right\rfloor.$$

Combining an idea of Babenko and Pichugov [2] with Theorem 13, Bojanov [3] obtained an explicit value of $B_{n,2}$.

Following Bojanov [3], let $P \in W_n$ and $t_1 \leq t_2 \leq \dots \leq t_n$ be the zeros of $t \mapsto P(t)$. Then we have

$$P'(t) = P(t)\sigma(t), \quad P''(t) = P'(t)\sigma(t) + P(t)\sigma'(t) \quad (P \in W_n),$$

where

$$\sigma(t) = \sum_{\nu=1}^n \frac{1}{t - t_k}.$$

Suppose that $\|P\|_{\infty} = |P(\tau)|$ and $\tau \in (-1, 1)$. Then $P'(\tau) = 0$ and therefore $\sigma(\tau) = 0$. Thus, $|P''(\tau)| = |\sigma'(\tau)|$. Choose $P = p_{n,k}$, where $k = 1, \dots, n-1$. Then $\tau = b_{n,k} = (2k - n)/n$ and

$$\sigma(t) = \frac{k}{t+1} + \frac{n-k}{t-1}.$$

Therefore,

$$\|p''_{n,k}\|_{\infty} \geq |p''_{n,k}(\tau)| = |\sigma'(\tau)| = \frac{n^2}{4} \left(\frac{1}{k} + \frac{1}{n-k} \right).$$

But the last expression attains its minimal value for $k = \lfloor n/2 \rfloor$ and this minimal value is n for even n , respectively $n(1 + 1/(n^2 - 1))$, for odd n . Adding the obvious fact that

$$\|p''_{n,1}\|_{\infty} = \|p''_{n,n}\|_{\infty} = \frac{1}{4} n(n-1) \geq n \quad (\text{for } n > 4),$$

we get $B_{n,2} = n$ for even $n \geq 6$, and

$$B_{n,2} \geq n \left(1 + \frac{1}{n^2 - 1} \right) \quad \text{for odd } n \geq 5.$$

Bojanov [3] also proved:

Theorem 14 *Let $x \mapsto \varphi(x)$ be any continuously differentiable, strictly increasing convex function in $[0, +\infty)$. Then for every $n \in \mathbb{N}$ and $m \in \{1, \dots, n\}$,*

$$\int_{-1}^1 \varphi(|P^{(m)}(t)|) dt \geq \int_{-1}^1 \varphi(|p_{n,n}^{(m)}(t)|) dt$$

for every polynomial $P \in W_n$ such that $P(1) = 1$.

If $\varphi(x) = x^r$ ($1 \leq r < +\infty$) this theorem reduces to the following result:

Corollary 15 *Let $P \in \mathcal{P}_n$, $P(1) = 1$, and $1 \leq r < +\infty$. Then*

$$\|P^{(m)}\|_r \geq \frac{n!}{2^m(n-m)!} \left(\frac{2}{(n-m)r+1} \right)^{1/r}.$$

Notice that for $m = 1$ this corollary gives Theorem 11.

Inequalities of Turán type for trigonometric polynomials were investigated by Babenko and Pichugov [1]–[2], Zhou [20], Tyrygin [12]–[13], and Bojanov [3]–[4].

4 A Result of Chen

In this section we mention a recent result of Chen [5], which can be expressed in the same way as the Markov's inequality in [7] (see also [9]).

We consider a general case with a given non-negative measure $d\sigma(t)$ on the real line $\mathbb{R}$, with compact or infinite support, for which all moments

$$\mu_\nu = \int_{\mathbb{R}} t^\nu d\sigma(t), \quad \nu = 0, 1, \dots,$$

exist and are finite, and $\mu_0 > 0$. Then there exists a unique set of orthonormal polynomials $\pi_\nu(\cdot) = \pi_\nu(\cdot; d\sigma)$, $\nu = 0, 1, \dots$, defined by

$$\pi_\nu(t) = a_\nu t^\nu + \text{lower degree terms}, \quad a_\nu > 0,$$

and

$$\int_{\mathbb{R}} \pi_\nu(t) \pi_\mu(t) d\sigma(t) = \delta_{\nu\mu}, \quad \nu, \mu \geq 0. \quad (8)$$

For each polynomial $P \in \mathcal{P}_n$, with complex coefficients, we take

$$\|P\| = \left(\int_{\mathbb{R}} |P(t)|^2 d\sigma(t) \right)^{1/2}.$$

As a restricted subset of $\mathcal{P}_n$ Chen [5] took

$$W_n = \mathcal{P}_{n,m}(d\sigma) = \{P \in \mathcal{P}_n \mid P \perp \mathcal{P}_{m-1}\},$$

i.e., $P \in W_n$ if $P \in \mathcal{P}_n$ and $(P, \pi_\nu) = 0$ for each $\nu = 0, 1, \dots, m-1$.

Consider now the extremal problem

$$A_{n,m} = A_{n,m}(d\sigma) = \inf_{P \in W_n} \frac{\|P^{(m)}\|}{\|P\|} \quad (1 \leq m \leq n). \quad (9)$$

Theorem 16 *The best constant $A_{n,m}$ defined in (9) is given by*

$$A_{n,m} = (\lambda_{\min}(B_{n,m}))^{1/2}, \quad (10)$$

where $\lambda_{\min}(B_{n,m})$ is the minimal eigenvalue of the matrix

$$B_{n,m} = [b_{i,j}^{(m)}]_{m \leq i,j \leq n},$$

whose elements are given by

$$b_{i,j}^{(m)} = \int_{\mathbb{R}} \pi_i^{(m)}(t) \pi_j^{(m)}(t) d\sigma(t), \quad m \leq i, j \leq n. \quad (11)$$

An extremal polynomial is

$$P^*(t) = \sum_{\nu=m}^n c_\nu \pi_\nu(t),$$

where $[c_k, c_{k+1}, \dots, c_n]^T$ is an eigenvector of the matrix $B_{n,m}$ corresponding to the eigenvalue $\lambda_{\min}(B_{n,m})$.

Proof. Let $P \in W_n$. Then we can write $P(t) = \sum_{\nu=m}^n c_\nu \pi_\nu(t)$ and

$$P^{(m)}(t) = \sum_{\nu=m}^n c_\nu \pi_\nu^{(m)}(t), \quad m \leq n,$$

where the coefficients c_ν are uniquely determined. Hence, by (8) and (11), we have

$$\|P\|^2 = \sum_{\nu=m}^n |c_\nu|^2 \quad \text{and} \quad \|P^{(m)}\|^2 = \sum_{i,j=m}^n c_i \bar{c}_j b_{i,j}^{(m)}.$$

Then

$$\frac{\|P^{(m)}\|^2}{\|P\|^2} = \frac{\sum_{i,j=m}^n c_i \bar{c}_j b_{i,j}^{(m)}}{\sum_{i=m}^n |c_i|^2} = \frac{\langle B_{n,m} \mathbf{c}, \mathbf{c} \rangle}{\langle \mathbf{c}, \mathbf{c} \rangle}, \quad (12)$$

where $\langle \cdot, \cdot \rangle$ denotes the standard inner product in an $(n-m+1)$ -dimensional space.

The matrix $B_{n,m}$ is evidently positive definite. Since the right side in (12) is not smaller than the minimal eigenvalue of this matrix, we obtain

$$\|P^{(m)}\|^2 \geq \lambda_{\min}(B_{n,m}) \|P\|^2. \quad (13)$$

In order to show that $A_{n,m}$, given by (10), is the best possible, we note that (13) reduces to an equality if we put $P(t) = P^*(t) = \sum_{\nu=m}^n c_\nu^* \pi_\nu(t)$, where $[c_m^*, c_{m+1}^*, \dots, c_n^*]^T$ is an eigenvector of the matrix $B_{n,m}$ corresponding to $\lambda_{\min}(B_{n,m})$. Q.E.D.

An alternative result like Theorem 16 is the following theorem:

Theorem 17 Let $Q_{n,m} = [q_{ij}^{(m)}]_{m \leq i,j \leq n}$ be an upper triangular matrix of the order $n-m+1$, whose elements $q_{ij}^{(m)}$ are given by the following inner product

$$q_{ij}^{(m)} = (\pi_j^{(m)}, \pi_{i-m}) \quad (m \leq i, j \leq n).$$

Then the best constant $A_{n,m}$ defined in (9) is given by

$$A_{n,m} = (\lambda_{\min}(Q_{n,m} Q_{n,m}^T))^{1/2}. \quad (14)$$

Alternatively, (14) can be expressed in the form

$$A_{n,m} = (\lambda_{\max}(C_{n,m}))^{-1/2}, \quad (15)$$

where $C_{n,m} = (Q_{n,m} Q_{n,m}^T)^{-1}$.

Proof. It is enough to consider only a real polynomial set $\mathcal{P}_n$. Let $P \in W_n$ and $\pi_j^{(m)}(t) = \sum_{i=m}^n q_{ij}^{(m)} \pi_{i-m}(t)$, where $q_{ij}^{(m)} = (\pi_j^{(m)}, \pi_{i-m})$. Then

$$P^{(m)}(t) = \sum_{j=m}^n c_j \sum_{i=m}^j q_{ij}^{(m)} \pi_{i-m}(t) = \sum_{i=m}^n \left(\sum_{j=m}^i c_j q_{ij}^{(m)} \right) \pi_{i-m}(t)$$

and

$$\|P^{(m)}\|^2 = \sum_{i=m}^n \left(\sum_{j=i}^n c_j q_{ij}^{(m)} \right)^2 = \sum_{i=m}^n y_i^2,$$

where

$$y_i = \sum_{j=i}^n c_j q_{ij}^{(m)}, \quad i = m, \dots, n. \quad (16)$$

Let $\mathbf{c} = [c_m, \dots, c_n]^T$, $\mathbf{y} = [y_m, \dots, y_n]^T$, and $Q_{n,m} = [q_{ij}^{(m)}]_{m \leq i, j \leq n}$. Since $\mathbf{y} = Q_{n,m} \mathbf{c}$, it follows that

$$\frac{\|P^{(m)}\|^2}{\|P\|^2} = \frac{\langle \mathbf{y}, \mathbf{y} \rangle}{\langle \mathbf{c}, \mathbf{c} \rangle} = \frac{\langle \mathbf{y}, \mathbf{y} \rangle}{\langle (Q_{n,m} Q_{n,m}^T)^{-1} \mathbf{y}, \mathbf{y} \rangle}.$$

Thus (14) and (15) hold. Q.E.D.

Now, we will consider a few special measures.

1° $d\sigma(t) = e^{-t^2} dt$, $-\infty < t < +\infty$. Here we have

$$\pi_\nu(t) = \hat{H}_\nu(t) = (\sqrt{\pi} 2^\nu \nu!)^{-1/2} H_\nu(t),$$

where H_ν is a Hermite polynomial of degree ν . Since

$$H'_\nu(t) = 2\nu H_{\nu-1}(t) \quad \text{and} \quad \hat{H}'_\nu(t) = \sqrt{2\nu} \hat{H}_{\nu-1}(t),$$

we have

$$\hat{H}_\nu^{(m)}(t) = \sqrt{2\nu} \sqrt{2(\nu-1)} \cdots \sqrt{2(\nu-m+1)} \hat{H}_{\nu-m}(t),$$

i.e.,

$$\hat{H}_\nu^{(m)}(t) = \sqrt{2^m m!} \binom{\nu}{m} \hat{H}_{\nu-m}(t),$$

and

$$b_{ij}^{(m)} = 2^m m! \binom{i}{m} \delta_{ij}, \quad m \leq i, j \leq n.$$

Thus, we find $\lambda_{\min}(B_{n,m}) = 2^m m!$ and $A_{n,m} = 2^{m/2} \sqrt{m!}$.

2° $d\sigma(t) = t^s e^{-t} dt$, $0 < t < +\infty$. Here we have the generalized Laguerre case with

$$\pi_\nu(t) = \hat{L}_\nu^s(t) = \sqrt{\nu!/\Gamma(\nu+s+1)} \sum_{i=0}^{\nu} (-1)^{\nu-i} \binom{\nu+s}{\nu-i} \frac{t^i}{i!},$$

where Γ is the gamma function.

First, we consider the simplest case where $m = 1$. Since

$$\frac{d}{dt} \hat{L}_j^s(t) = \sum_{i=1}^j q_{ij}^{(1)} \hat{L}_{i-1}^s(t), \quad q_{ij}^{(1)} = -\sqrt{\frac{j!}{\Gamma(j+s+1)}} \cdot \sqrt{\frac{\Gamma(i+s)}{(i-1)!}},$$

from the equalities (16), it follows that

$$c_i = y_{i+1} - \sqrt{\frac{i+s}{i}} y_i, \quad i = 1, \dots, n,$$

where we put $y_{n+1} = 0$. The elements $p_{ij}^{(1)}$ of the matrix $P_{n,1} = Q_{n,1}^{-1}$ are

$$p_{ij}^{(1)} = -\sqrt{1 + \frac{s}{i}}, \quad i = 1, \dots, n; \quad p_{i,i+1}^{(1)} = 1, \quad i = 1, \dots, n-1;$$

$$p_{ij}^{(1)} = 0, \quad \text{otherwise,}$$

so that $C_{n,1} = P_{n,1}^T P_{n,1} = -J_n$, where

$$J_n = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \mathbf{0} \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \alpha_2 & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_{n-1}} \\ \mathbf{0} & & & \sqrt{\beta_{n-1}} & \alpha_{n-1} \end{bmatrix}$$

and

$$\alpha_0 = -(1+s), \quad \alpha_\nu = -\left(2 + \frac{s}{\nu+1}\right), \quad \beta_\nu = 1 + \frac{s}{\nu}, \quad \nu = 1, \dots, n-1.$$

We see that J_n is the Jacobi matrix for monic orthogonal polynomials $\{Q_\nu\}$, which satisfy the following three-term recurrence relation

$$Q_{\nu+1}(t) = (t - \alpha_\nu)Q_\nu(t) - \beta_\nu Q_{\nu-1}(t), \quad \nu = 0, 1, \dots,$$

with $Q_{-1}(t) = 0$ and $Q_0(t) = 1$. The eigenvalues of $C_{n,1}$ are $\lambda_\nu = -t_\nu$, where $Q_n(t_\nu) = 0$ for $\nu = 1, \dots, n$.

The standard Laguerre case ($s = 0$) can be exactly solved. In fact, for $t = 2(z-1)$ with $-1 \leq z \leq 1$, we have

$$Q_\nu(t) = \cos(2\nu+1)\frac{\theta}{2} \Big/ \cos \frac{\theta}{2}, \quad z = \cos \theta.$$

The eigenvalues of the matrix $C_{n,1}$ are

$$\lambda_\nu = -t_\nu = 4 \sin^2 \frac{(2\nu - 1)\pi}{2(2n + 1)}, \quad \nu = 1, \dots, n.$$

Since $\lambda_{\max}(C_{n,1}) = \lambda_n$, we obtain

$$A_{n,1} = \left(2 \cos \frac{\pi}{2n + 1} \right)^{-1}.$$

Now, we consider the case when $m = 2$ and $s = 0$. First, we note that

$$\frac{d^m}{dt^m} \hat{L}_j(t) = (-1)^m \sum_{i=m}^j \binom{j-i+m-1}{m-1} \hat{L}_{i-m}(t).$$

The formulae (16), for $m = 2$, become

$$y_i = \sum_{j=i}^n (j - i + 1) c_j, \quad i = 2, \dots, n.$$

Since $\Delta^2 y_i = c_i$ ($y_{n+1} = y_{n+2} = 0$), we find a five-diagonal symmetric matrix of order $n - 1$

$$C_{n,2} = \begin{bmatrix} 1 & -2 & 1 & & & & & \mathbf{O} \\ -2 & 5 & -4 & 1 & & & & \\ & 1 & -4 & 6 & -4 & 1 & & \\ & & 1 & -4 & 6 & -4 & 1 & \\ & & & \ddots & \ddots & \ddots & \ddots & \ddots \\ & & & & 1 & -4 & 6 & -4 & 1 \\ & & & & & 1 & -4 & 6 & -4 \\ \mathbf{O} & & & & & & 1 & -4 & 6 \end{bmatrix}.$$

Thus, using the maximal eigenvalue of this matrix, we obtain the best constant $A_{n,2} = (\lambda_{\max}(C_{n,2}))^{-1/2}$. In the simplest case when $n = 2$ and $n = 3$ we have $A_{2,2} = 1$ and $A_{3,2} = (3 - 2\sqrt{2})^{1/2}$, respectively.

We conclude this paper with a remark that Varma [17] also studied an extremal problem on $(0, +\infty)$ with respect to the Laguerre measure, i.e., when $\|f\|_2^2 = \int_0^\infty e^{-t} f(t)^2 dt$.

Theorem 18 *Let P be an algebraic polynomial of degree n whose zeros τ_ν ($\nu = 1, \dots, n$) all lie in the interval $[0, \infty)$. If $P(0) = 0$ or*

$$\sum_{\nu=1}^n \tau_\nu^{-1} \geq \frac{1}{2};$$

then

$$\|P'\|_2^2 \geq \frac{n}{2(2n-1)} \|P\|_2^2.$$

The equality holds for $P(t) = t^n$.

References

- [1] V. F. Babenko and S. A. Pichugov, Exact inequality for the derivative of a trigonometric polynomial having only real zeros, *Mat. Zametki* **39** (1986), 330–336 (Russian).
- [2] V. F. Babenko and S. A. Pichugov, On inequality for the derivative of a polynomial with real zeros, *Ukrain. Mat. Zh.* **38** (1986), 411–416 (Russian).
- [3] B. D. Bojanov, Polynomial inequalities, in “Open Problems in Approximation Theory” (B. Bojanov, Ed.), pp. 25–42, SCT, Singapore, 1993.
- [4] B. Bojanov, Turán's inequalities for trigonometric polynomials, *J. London. Math. Soc.* (2) **53** (1996), 539–550.
- [5] W. Chen, On the converse inequality of Markov type in L_2 norm, *J. Math. Anal. Appl.* **200** (1996), 708–716.
- [6] J. Erőd, Bizonyos polinomok maximumának, *Mat. Fiz. Lapok* **46** (1939), 58–82.
- [7] G. V. Milovanović, Various extremal problems of Markov's type for algebraic polynomials, *Facta Univ. Ser. Math. Inform.* **2** (1987), 7–28.
- [8] G. V. Milovanović, D. S. Mitrinović and Th. M. Rassias, On some Turán's extremal problems for algebraic polynomials, in “Topics in Polynomials of One and Several Variables and Their Applications: A Mathematical Legacy of P. L. Chebyshev (1821–1894)” (Th. M. Rassias, H. M. Srivastava, A. Yanushauskas, Eds.), pp. 403–433, World Scientific, Singapore, 1993.

- [9] G. V. Milovanović, D. S. Mitrinović and Th. M. Rassias, "Topics in Polynomials: Extremal Problems, Inequalities, Zeros", World Scientific, Singapore – New Jersey – London – Hong Kong, 1994.
- [10] J. Szabados and A. K. Varma, Inequalities for derivatives of polynomial having real zeros, in "Approximation Theory III" (E. W. Cheney, Ed.), pp. 881–887, Academic Press, New York, 1980.
- [11] P. Turán, Über die Ableitung von Polynomen, *Composito Math.* **7** (1939), 85–95.
- [12] I. Ya. Tyrygin, Inequalities of P. Turán type in mixed integral metrics, *Dokl. Akad. Nauk Ukrain. SSR Ser. A* **1988**, no. 9, 14–17 (Russian).
- [13] I. Ya. Tyrygin, On inequalities of Turán type in some integral metrics, *Ukrain. Mat. Zh.* **40** (1988), 256–260 (Russian).
- [14] B. Underhill and A. K. Varma, An extension of some inequalities of P. Erdős and P. Turán concerning algebraic polynomials, *Acta Math. Hung.* **73** (1996), 1–28.
- [15] A. K. Varma, An analogue of some inequalities of P. Turán concerning algebraic polynomials having all zeros inside $[-1, 1]$, *Proc. Amer. Math. Soc.* **55** (1976), 305–309.
- [16] A. K. Varma, An analogue of some inequalities of P. Turán concerning algebraic polynomials having all zeros inside $[-1, 1]$. II, *Proc. Amer. Math. Soc.* **69** (1978), 25–33.
- [17] A. K. Varma, Some inequalities of algebraic polynomials having real zeros, *Proc. Amer. Math. Soc.* **75** (1979), 243–250.
- [18] A. K. Varma, Some inequalities of algebraic polynomials having all zeros inside $[-1, 1]$, *Proc. Amer. Math. Soc.* **88** (1983), 227–233.
- [19] A. K. Varma, Some inequalities of algebraic polynomials, *Proc. Amer. Math. Soc.* **123** (1995), 2041–2048.
- [20] S. P. Zhou, On Turán inequality in L_p -norm, *J. Hangzhou Univ.* **11** (1984), 28–33 (Chinese, English summary).
- [21] S. P. Zhou, An extension of the Turán inequality in L^p for $0 < p < 1$, *J. Math. Res. Exposition* **6**, No. 2 (1986), 27–30.
- [22] S. P. Zhou, Some remarks on Turán inequality, *J. Approx. Theory* **68** (1992), 45–48.